

Quasi-point separation of variables for the Henon-Heiles system and a system with quartic potential.

S. Rauch-Wojciechowski

Department of Mathematics, Linköping University, S-58183, Linköping, Sweden

A.V. Tsiganov

Department of Mathematical and Computational Physics, Institute of Physics,

St.Petersburg University, 198 904, St.Petersburg, Russia

Abstract

We examine the problem of integrability of two-dimensional Hamiltonian systems by means of separation of variables. The systematic approach to construction of the special non-pure coordinate separation of variables for certain natural two-dimensional Hamiltonians is proposed.

1 Introduction

In this note we study quasi-point separation of variables for certain natural Hamiltonians

$$H = \frac{1}{2}(p_x^2 + p_y^2) + V(x, y) \quad (1.1)$$

of two degrees of freedom.

The classical separability theory [10, 4] is concerned with orthogonal point transformations $x = \Phi_1(u, v)$ and $y = \Phi_2(u, v)$ for which the corresponding Hamilton-Jacobi equation expressed in terms of (u, v) -variables can be solved by separation of variables. The list of coordinate systems in plane that provide point separation of variables for systems with the Hamiltonian (1.1) are well known. Even an effective criterion of separability for a given potential $V(x, y)$ has been formulated [5, 10]. In this note we consider a non-pure point transformation related to a cartesian system of coordinates.

Very little is known about general canonical transformations $x_k = \Phi_k(u_j, p_{u_j})$ and $p_k = \Psi_k(u_j, p_{u_j})$, $k = 1, 2$, which separate the Hamiltonian (1.1). Recently, a special type of non-point transformations $x = \varphi(H, C) \cdot \Phi(u, v)$ has been introduced for the third integrable case of the Henon-Heiles system [7, 3] and for a system with a quartic potential [8]. Here $\varphi(H, C)$ is a certain function of the Hamiltonian H and of the second integral of motion C . On the orbit $\mathcal{O} (H = \alpha_1 \quad C = \alpha_2)$ this transformation becomes a point transformation and, therefore, we shall call it a quasi-point transformation. These transformations have been found either in the context of the Painlevé expansion [7, 8] or

have been derived from the Miura transformation for the systems related to stationary flows of soliton equations [3, 2]. However no general principle of constructing general (this means non point) canonically separable potentials is known. This is the reason why in this note we address to this question directly by starting with general separated equations and by trying to find transformations which lead to natural Hamiltonians (1.1) of two degrees of freedom. We do not find any essentially new potentials. We present in detail our construction of separating variables and, then, list the main results. We explain the connection of quasi-point transformations with the supersymmetric quantum mechanics.

2 Quasi-point canonical transformation

Let us begin with a two-dimensional system expressed in terms of canonical variables of separation (u, p_u, v, p_v) with the rectangular separated equations of the general form [4, 10]

$$\begin{aligned}\Delta_1(u, p_u) &= f(u)p_u^2 + V_1(u), \\ \Delta_2(v, p_v) &= g(v)p_v^2 + V_2(v).\end{aligned}\tag{2.1}$$

The functions Δ_j Poisson commute $\{\Delta_1, \Delta_2\} = 0$ with respect to the standard Poisson brackets. Notice that we could, by the using a point transformation, reduce these equations to equations with $f(u) = g(v) = 1$, but the form (2.1) is more suitable for further calculations.

As commuting integrals of motion we choose

$$H = \Delta_1 + \Delta_2 = f(u)p_u^2 + g(v)p_v^2 + V_+(u, v),\tag{2.2}$$

$$C = a\Delta_1 - b\Delta_2 = a f(u)p_u^2 - b g(v)p_v^2 + V_-(u, v), \quad a, b \in \mathbb{R}$$

which clearly define $V_{\pm}(u, v)$. These integrals are second order polynomials in momenta. A condition of separability for the potentials $V_{\pm}$ is

$$\frac{\partial^2}{\partial u \partial v} V_{\pm}(u, v) = 0.\tag{2.3}$$

Let us denote a canonical transformation to the cartesian variables (x, y) as

$$x = \Phi_1(u, p_u, v, p_v) \quad \text{and} \quad y = \Phi_2(u, p_u, v, p_v),\tag{2.4}$$

and let us require that after the transformation (2.4) the Hamiltonian (2.2) takes the natural Hamiltonian form (1.1). Then

$$p_x = \{H, x\} \quad \text{and} \quad p_y = \{H, y\}$$

and, after substituting these momenta into the canonical Poisson brackets, we shall get a system of equations for the transformations (2.4). It reads

$$\begin{aligned}\{\Phi_j(u, p_u, v, p_v), \Phi_k(u, p_u, v, p_v)\} &= 0, \\ \{\{H, \Phi_j(u, p_u, v, p_v)\}, \{H, \Phi_k(u, p_u, v, p_v)\}\} &= 0, \\ \{\{H, \Phi_j(u, p_u, v, p_v)\}, \Phi_k(u, p_u, v, p_v)\} &= \delta_{jk}, \\ &(j, k = 1, 2).\end{aligned}\tag{2.5}$$

Since, we can not solve this system of equations in the whole generality, we have to make certain simplifying ansatz for the functions Φ_k in order to obtain certain particular solutions of (2.5).

In the class of point transformations $\tilde{x} = \tilde{\Phi}_1(u, v)$ and $\tilde{y} = \tilde{\Phi}_2(u, v)$ the general solution of (2.5) has the form

$$\tilde{x} = \alpha u + \beta v + \lambda_1, \quad \tilde{y} = \gamma u + \delta v + \lambda_2, \quad (2.6)$$

which is a superposition of rotation and translation transformations. Since the functions $f(u)$, $g(v)$ and $V_{\pm}(u, v)$ are arbitrary we can always rewrite the solution (2.6) in the following nonsymmetric form

$$\tilde{x} = \tilde{\Phi}_1(u, v) = u - v, \quad \tilde{y} = \tilde{\Phi}_2(u, v) = u + v + 2\gamma, \quad (2.7)$$

which will be used below. The polynomial order of integrals of motion (2.2) remains unchanged under (2.7).

In order to obtain a non-point transformation as a solution of (2.5) we shall introduce a particular ansatz for $x = \Phi_1(u, p_u, v, p_v)$, which is suggested by the results of [7, 8]. For explaining the structure of this ansatz we begin with a simple example, which will be present in detail below. Let us take

$$\Delta_j(\xi) = \xi^n p_{\xi}^2 + V_j(\xi), \quad \xi = u, v, \quad (2.8)$$

$$x^2 = \Delta_1(u)u^{-m}, \quad u^m = \Delta_1x^{-2},$$

and require that the Hamiltonian (2.2) takes the natural form (1.1), when expressed through the new variables (x, p_x) . Then, we obtain

$$\frac{p_x}{x} = -mu^{n-1}p_u$$

and by applying the Poisson brackets (2.5) we get

$$\left\{ \frac{p_x}{x}, x^2 \right\} = 2 = \left\{ -mu^{n-1}p_u, u^{n-m}p_u^2 + u^{-m}V_1(u) \right\}. \quad (2.9)$$

The kinetic part of this equation leads to a restriction for the kinetic part of the Hamiltonian: $n = 2 - m$. The potential part of (2.9) gives

$$u \frac{\partial V_1(u)}{\partial u} - mV_1(u) = -2m^{-1}u^{2m},$$

which has the solution

$$V_1 = -2m^{-2}u^{2m} + \alpha u^m = -2m^{-2}\Delta_1^2x^{-4} + \alpha\Delta_1x^2, \quad \alpha \in \mathbb{R}. \quad (2.10)$$

According to the definitions (1.1) and (2.8) the momenta p_u is

$$p_u = -\frac{\Delta_1 p_x x^{-3}}{mu},$$

and, after substituting u, p_u and $V_1(u)$, the separated equation Δ_1 (2.8) becomes

$$\Delta_1(x, p_x) = \frac{p_x^2}{2} - \frac{m^2}{2}(x^4 - \alpha x^2), \quad \alpha \in \mathbb{R}.$$

The second pair of variables (y, p_y) follows from the second separated equation $\Delta_2(v)$ and the Hamiltonian (2.2) transforms to the new Hamiltonian

$$H = \frac{1}{2}(p_x^2 + p_y^2) + \alpha_1 x^4 + \alpha_2 y^4 + \alpha_3 x^2 + \alpha_4 y^2, \quad \alpha_k \in \mathbb{R},$$

under non-point canonical transformation (2.8).

Next we consider a symmetric form of this transformation related to the symmetric form of the second integral of motion (2.2). Let us take

$$P_n(x) = \sum_{k=0}^n a_k x^k = C \phi(u - v), \quad a_k \in \mathbb{R}, \quad (2.11)$$

where $P_n(x)$ is a n -th order polynomial, C is an integral of motion (2.2) and $\phi(u - v)$ is a function of one variable $z = u - v$. On the orbit $\mathcal{O}$ ($C = c = \text{const}$) the transformation (2.11) becomes a point transformation and, therefore, we shall call this transformation a quasi-point transformation.

After substituting the ansatz (2.11) into the equation

$$\{\{H(u, p_u, v, p_v), \Phi_1(u, p_u, v, p_v)\}, \Phi_1(u, p_u, v, p_v)\} = 1 \quad (2.12)$$

we consider terms at independent powers of momenta. It appears that a solution of the corresponding equation exist only if $P_n(x) = x^2$ (up to a point transformation (2.6)) and we get one equation for the function $\phi(z)$

$$2 \frac{d\phi(z)}{dz} = \phi(z) \frac{d^2\phi(z)}{dz^2}, \quad z = u - v, \quad (2.13)$$

and a system of equations for the functions $f(u)$ and $g(v)$

$$\begin{aligned} f(u) - bg(v) - b(u - v) \frac{dg(v)}{dv} &= 0, \\ -af(u) + bg(v) + a(u - v) \frac{df(u)}{du} &= 0. \end{aligned} \quad (2.14)$$

The general solutions for (2.13 and 2.14) are

$$\begin{aligned} \phi(z) &= z^{-1} = \tilde{\Phi}_1^{-1}(u, v), \\ a = b &= \alpha_1, \quad f(u) = \alpha_2 u + \alpha_3, \quad g(v) = \alpha_2 v + \alpha_3, \quad \alpha_k \in \mathbb{R}. \end{aligned} \quad (2.15)$$

Further, we obtain that

$$x^2 = C \tilde{\Phi}_1^{-1}(u, v),$$

where $\tilde{\Phi}_1^{-1}(u, v)$ is a point transformation (2.7) and, therefore, we can consider (2.11) as a natural generalization of a pure point transformation, which can be applied to the

other three types of coordinate separated equations in the plain. For instance for the parabolic system of coordinates we have to substitute

$$C = \frac{v^2 \Delta_1 - u^2 \Delta_2}{u^2 + v^2}, \quad \tilde{\Phi}_1(u, v) = u^2 - v^2,$$

into the ansatz (2.11).

From the remaining terms of (2.12) (by taking into account (2.13) and (2.14)) we obtain that the potential $V_-(u, v)$ obeys the following equation

$$\left((\alpha_2 u + \alpha_3) \frac{\partial V_-}{\partial u} - (\alpha_2 v + \alpha_3) \frac{\partial V_-}{\partial v} \right) - \frac{(\alpha_2(u+v) + 2\alpha_3)}{\alpha_1(u-v)} V_- = -\frac{2(u-v)^3}{\alpha_1(u-v)}. \quad (2.16)$$

For a separable potential $\partial^2 V_-(u, v)/\partial u \partial v = 0$ this equation has a unique solution

$$V_{1,2}(\xi) = \frac{2/\alpha_1 \xi^3 + \beta_2 \xi^2 + \beta_1 \xi + \beta_0}{\alpha_2 \xi + \alpha_3}; \quad \xi = u, v. \quad (2.17)$$

where β_k are arbitrary constants.

Next we have to determine the second variable $y = \Phi_2(u, p_u, v, p_v)$ by using equations (2.5). Again we are not able to find a general solution for Φ_2 and we shall use the particular ansatz of [7, 8]

$$Q_n(y) = \sum_{k=0}^n b_k y^k = \psi_1(u+v) + \psi_2(x, p_x), \quad p_y = \{H, y\}, \quad (2.18)$$

where $Q_n(y)$ is a polynomial of the order n and ψ_k are unspecified as yet functions.

After substituting (2.18) into the equations

$$\{x, Q_n(y)\} = 0 \quad \text{and} \quad \{p_x, Q_n(y)\} = 0$$

(recall that $p_x = \{H, x\}$) we obtain

$$Q_n(y) = \gamma_1 \left[u + v + \gamma_2 - \frac{\alpha_1}{2} \left(\frac{p_x}{x} \right)^2 + \frac{\alpha_2 x^2}{2} \right]. \quad (2.19)$$

This ansatz is a superposition of a point transformation $\tilde{\Phi}_2(u, v)$ (2.7) with a term depending on the first variables (x, p_x) .

By using the remaining equation $\{p_y, y\} = 1$ we obtain (up to a point transformations (2.6)) that

$$Q_n(y) = y^2 \quad \gamma_1 = -\frac{1}{\alpha_2}, \quad (2.20)$$

and yet another differential equation of the second order for the potential $V_+(u, v)$. We present here a partial form of this equation with $\alpha_2 = 0$

$$\frac{\partial^2 V_+}{\partial z^2} = z \frac{\partial V_+}{\partial z}, \quad z = u - v.$$

This equation for V_+ is equivalent to the equation for V_- (2.16) provided that the constants $\alpha_1, \alpha_2, \beta_2$ and γ_2 satisfy

$$\gamma_2 + \alpha_1 \beta_2 + 2\alpha_3 = 0. \quad (2.21)$$

Let us summarize these considerations as

Proposition 1 *A quasi-point transformation of the form*

$$\begin{aligned} P_n(x) &= \sum_{k=1}^n a_k x^k = \frac{C}{u-v}, \\ Q_m(y) &= \sum_{k=1}^m b_k y^k = \psi_1(u+v) + \psi_2(x, p_x) \quad a_k, b_k \in \mathbb{R} \end{aligned}$$

extended to a canonical transformation, transforms the Hamiltonian

$$H = f(u)p_u^2 + g(v)p_v^2 + V_1(u) + V_2(v)$$

into the natural Hamiltonian form

$$H = \frac{1}{2}(p_x^2 + p_y^2) + V(x, y),$$

if and only if

$$\begin{aligned} x^2 &= \frac{C}{u-v}, \\ \frac{p_x}{x} &= -\frac{(\alpha_2 u + \alpha_3)p_u - (\alpha_2 v + \alpha_3)p_v}{u-v} \\ y^2 &= -\frac{1}{\alpha_2} \left(u + v - \alpha_1 \beta_2 - 2\alpha_3 - \frac{\alpha_1}{2} \left(\frac{p_x}{x} \right)^2 + \frac{\alpha_2 x^2}{2} \right), \\ y p_y &= \frac{1}{\alpha_2} \left((\alpha_2 u + \alpha_3)p_u - (\alpha_2 v + \alpha_3)p_v \right. \\ &\quad \left. + \frac{p_x}{2x} \left(\frac{\alpha_1 p_x^2}{x^2} + \frac{6\alpha_3}{\alpha_2} - 6\alpha_2 y^2 + \alpha_1 \beta_2 - \alpha_2 x^2 \right) \right). \end{aligned} \tag{2.22}$$

and the potentials

$$V_k(\xi) = \frac{2/\alpha_1 \xi^3 + \beta_2 \xi^2 + \beta_1 \xi + \beta_0}{\alpha_2 \xi + \alpha_3}; \quad k = 1, \xi = u \quad \text{or} \quad k = 2, \xi = v.$$

Constants $\alpha_j, \beta_j, j = 1, 2, 3$ are a six free parameters of this transformation.

An inverse transformation to (2.22) reads

$$\begin{aligned} \xi &= \pm \frac{C}{2x^2} + \frac{\alpha_1 p_x^2}{4x^2} - \frac{\alpha_2}{4}(2y^2 + x^2) - \gamma_2; \quad \xi = u, v, \\ p_\xi &= \frac{1}{2(\alpha_2 \xi + \alpha_3)} \left(\frac{p_x C}{x^3} - \alpha_2 y p_y - \right. \\ &\quad \left. - \frac{p_x}{2x} \left(\frac{\alpha_1 p_x^2}{x^2} + \frac{6\alpha_3}{\alpha_2} - 6\alpha_2 y^2 + \alpha_1 \beta_2 - \alpha_2 x^2 \right) \right), \end{aligned} \tag{2.23}$$

where the integral C is an unspecified as yet function of the new variables (x, p_x, y, p_y) . We have to substitute this inverse transformation into the definition of the integral of motion C (2.2) and to solve the resulting equation. It has the polynomial form

$$aC^4 + bC^2 + c = 0, \quad (2.24)$$

with the coefficients a, b and c depending on the variables (x, p_x, y, p_y) . There are two solutions $C_{1,2}^2$, which are related by the change of variables $y \rightarrow -y$, $p_y \rightarrow -p_y$, and they are polynomial expressions in the new variables. Notice that the second integral of motion C (2.2), which is polynomial in (u, p_u, v, p_v) becomes an algebraic function of (x, p_x, y, p_y) .

After substitution of $C(x, p_x, y, p_y)$ into (2.23) we get an inverse transformation.

Proposition 2 *A Hamiltonian*

$$H = f(u)p_u^2 + g(v)p_v^2 + V_1(u) + V_2(v)$$

with the potentials given by (2.17) transforms to either of two natural Hamiltonians

$$\begin{aligned} H_1 &= \frac{1}{2}(p_x^2 + p_y^2) - \frac{\alpha_2}{2\alpha_1}(x^4 + 6x^2y^3 + 8y^4) \\ &+ \frac{\beta_2}{2}(x^2 + 4y^2) + \frac{2\beta_0}{\alpha_2^2 y^2} + 2\alpha_2\beta_1, \end{aligned} \quad (2.25)$$

and

$$H_2 = \frac{1}{2}(p_x^2 + p_y^2) - \frac{\alpha_2}{2\alpha_1}(x^4 + 6x^2y^3 + y^4), \quad (2.26)$$

under the quasi-point transformation (2.22) and under the point transformation, respectively.

For brevity, we put $\alpha_3 = 0$ in H_1 (2.25), since the introduction of $\alpha_3 \neq 0$ corresponds to the shift of the variable $y \rightarrow y + \alpha_3/4\alpha_2$ [9]. In H_2 (2.26) we presented only the highest polynomial terms.

Systems with the Hamiltonians H_1 and H_2 are associated with restricted flows of some PDE's [3, 2]. In classical mechanics these systems have common separated equations. Second integrals of motion C^2 derived from the equation (2.24) are well known and can be founded in [8, 9].

By rescaling constants a_j, β_j and by taking the limits $\alpha_2 \rightarrow 0$ and $\alpha_3 \rightarrow 1$, the Hamiltonians (2.25) and (2.26) are transformed to the following Hamiltonians for the Henon-Heiles system [3, 7]

$$\begin{aligned} H_1 &= \frac{1}{2}(p_x^2 + p_y^2) + \frac{2}{\alpha_1\alpha_3}y(3x^2 + 16y^2) - \beta_2(x^2 + 16y^2) \\ &+ 2(\alpha_1\beta_2 - 2\beta_1)y + 2\beta_0 + \alpha_1\beta_2\beta_1, \\ H_2 &= \frac{p_x^2 + p_y^2}{2} + ay(x^2 + 2y^2). \end{aligned} \quad (2.27)$$

A complete account of this limit procedure can be found in [9].

It is known that the Hamiltonian (2.25) has an integrable extensions

$$H = H_{1,2} + \frac{m}{x^2} + \frac{l}{y^2} + \frac{n}{x^6} + ey,$$

where either $e = 0$, or $n = m = 0$ [8]. We can include the terms μx^{-2} and ly^{-2} into our proposed scheme. Let us consider the following infinite-dimensional representation of $sl(2)$ defined in Cartan-Weil basis

$$s_3 = \frac{xp}{2}, \quad s_+ = \frac{x^2}{2}, \quad s_- = -\frac{p^2}{2}, \quad \{p, q\} = 1.$$

The mapping

$$s_3 \rightarrow s'_3 = s_3, \quad s_+ \rightarrow s'_+ = s_+, \quad (2.28)$$

$$s_- \rightarrow s'_- = s_- + fs_+^{-1} = \frac{p^2}{2} + \frac{2f}{x^2}, \quad f \in \mathbb{R}$$

is an outer automorphism of the space of infinite-dimensional representations of $sl(2)$. For the Henon-Heiles system and for the system with quartic potential the phase space can be identified completely or partially with the coadjoint orbits in $sl(2)^*$ as (2.28). Hence, all the presented results can be carried over on the systems with shifted squared momenta. The corresponding deformation of the separated equation has been described in [8].

3 Quasi-point transformations and SUSY quantum mechanics

Next, we present the interesting relations of the quasi-point canonical transformation with the supersymmetrical quantum mechanics, which represents in a concise algebraic form the spectral equivalence between different Hamiltonian quantum systems realized by the Darboux transformation [1]. Further, for brevity, we fix the value of parameters $\beta_k = 0$, $\alpha_2 = 1$, $\alpha_1 = 2$.

Variables of separation u, v for the Hamiltonian H_2 can be defined as the roots of quadratic equation

$$\xi^2 - y\xi + \frac{y^2 - x^2}{4} = 0. \quad (3.29)$$

The quasi-point transformations (2.23) can be presented in similar form. Variables u, v (2.23) are roots of the quadratic equation

$$\xi^2 - \frac{(q_+ + q_-)}{x^2}\xi + \frac{(q_+ - q_-)^2}{4x^2} = 0. \quad (3.30)$$

Here, we introduced functions $q_{\pm}$ and f with the following properties

$$\{H, q_{\pm}\} = \pm f q_{\pm}, \quad C^2 = 4q_+ q_-. \quad (3.31)$$

This algebra (3.31) is the classical limit of the two-dimensional quantum SUSY algebra [1]. For the system with quartic potential functions $q_{\pm}$ and f are given by

$$\begin{aligned} q_+ &= \frac{1}{2}p_x^2 - \frac{1}{4}x^2(2y^2 + x^2) + x(y p_x - \frac{1}{2}p_y x), \\ q_-(x, p_x, y, p_y) &= q_+(x, p_x, -y, -p_y), \end{aligned} \quad (3.32)$$

$$f = 2y,$$

and for the Henon-Heiles system

$$\begin{aligned} q_+ &= \frac{1}{2}p_x^2 + 2x^2y + ix \cdot \sqrt{2(2p_x y - x p_y)p_x - (8y^2 + x^2)x^2}, \\ q_-(x, p_x, y, p_y) &= q_+^*(x, p_x, y, p_y), \end{aligned} \quad (3.33)$$

$$f = i \frac{4p_x y - x p_y}{\sqrt{2(2p_x y - x p_y)p_x - (8y^2 + x^2)x^2}}$$

where i is the imaginary unit and q_+^* means the complex conjugate of the function q_+ . For the arbitrary values of the parameters α_k and β_k these functions are derived by equations (2.23) and (3.30).

For the system with quartic potentials functions $q_{\pm}$ and f (3.32) have been introduced in [6] by considering a linearization of the corresponding Hamiltonian flow on certain constraint submanifolds.

It would be interesting to apply equation (3.30) and their quantum counterpart for separating variables in quantum mechanics.

Motivated by relation of classical limit of SUSY quantum mechanics and quasi-point separation of variables we present the classical SUSY algebra (3.31) for another two-dimensional natural Hamiltonian systems. For the systems of [11] and for the Holt-like systems the classical limit of SUSY algebra (3.31) is equal to

$$\begin{aligned} H_W &= \frac{p_x^2 + p_y^2}{2} + \left(\frac{a_1^2}{2} + \alpha_2 \alpha_3 \right) x^{2/3} - \frac{9}{16} \alpha_1^2 \alpha_2 x^{-2/3} y - \alpha_3 y, \\ q_+ &= -\frac{p_x^2}{2} + \frac{9}{16} \left(\alpha_1^2 \alpha_2 x^{-2/3} y - \frac{\alpha_2}{\alpha_1} \right) + \left(\frac{\alpha_1^2}{2} + \alpha_2 \alpha_3 \right) x^{2/3} + i \left(\alpha_1 x^{1/3} p_x + \frac{\alpha_2}{3\alpha_1} p_y \right), \\ f &= -\frac{2}{3} i \alpha_1 x^{-2/3} \end{aligned}$$

and

$$\begin{aligned} H_H &= \frac{p_x^2 + p_y^2}{2} + y^{-2/3} \left(\frac{9}{2} y^2 + x^2 \right), \\ q_+ &= p_x^2 + 2y^{-2/3} x^2 + i \cdot \sqrt{2(p_x p_y + 6x y^{1/3})^2 - (2x^2 y^{-2/3})^2}, \\ f &= \frac{4i x^2 p_y}{3y \sqrt{2(y^{2/3} p_x p_y + 6y x)^2 - 4x^4}}, \end{aligned}$$

where $q_-(x, p_x, y, p_y) = q_+^*(x, p_x, y, p_y)$.

Notice, that the construction of isospectral two-dimensional Hamiltonians in supersymmetrical quantum mechanics is closely connected with another problem, namely with a search for the second integral of motion for the quantum integrable systems [1]. Here we present a new characterisation of this problem.

Proposition 3 *Let us start with the classical SUSY algebra (3.31) for a two-dimensional integrable system defined by four functions H, f and $q^\pm$. If the following equation for Δh can be solved*

$$\{\Delta h, \{q_+, q_-\}\} = \{f, q_+\}q_- + \{f, q_-\}q_+, \quad (3.34)$$

then the pair of the mutually commuting integrals of motion

$$\widetilde{H} = H + \Delta h, \quad \widetilde{C} = \{q^+, q^-\}, \quad \{\widetilde{H}, \widetilde{C}\} = 0, \quad (3.35)$$

defines a new two-dimensional integrable system.

The evolution on a $2n$ -dimensional symplectic manifold is called completely integrable if there exists n functions $I_1, \dots, I_n$, which are independent integrals in the involution

$$\{I_i(x, p), I_j(x, p)\} = 0, \quad i, j = 1, \dots, n.$$

The initial integrals of motion H and C are functionally independent functions and, therefore, the new integrals $\widetilde{H}$ and $\widetilde{C}$ are independent. For the proof of involution of integrals we need the the Jacobi identity for the Poisson brackets. Assuming first (3.31) holds we have

$$\begin{aligned} \{H, \widetilde{C}\} &= \{H, \{q^+, q^-\}\} = \{q^+, \{H, q^-\}\} - \{q^-, \{H, q^+\}\} = \\ &= q^-\{f, q^+\} + q^+\{f, q^-\}. \end{aligned}$$

Using next (3.34) we get (3.35)

$$\{H + \Delta h, \widetilde{C}\} = \{\widetilde{H}, \widetilde{C}\} = 0.$$

So, we have two independent integrals of motion in involution, which are defined new integrable system.

As an example, for the system with quartic potential equation (3.34) can be integrated and it yields

$$\begin{aligned} H_1 &= \frac{p_x^2 + p_y^2}{2} - \frac{1}{4}(x^4 + 6x^2y^2 + 8y^4), \\ \Delta h &= -\frac{1}{8}x^4, \end{aligned} \quad (3.36)$$

$$\begin{aligned} \widetilde{H} &= H_1 + \Delta h = \frac{p_x^2 + p_y^2}{2} - \frac{1}{8}(x^4 + 12x^2y^2 + 16y^4), \\ \widetilde{C} &= \{q_+, q_-\} = 16(2(p_yx - yp_x)p_x - (2y^2 - x^2)yx^2). \end{aligned}$$

New system with integrals of motion $\widetilde{H}, \widetilde{C}$ is separable in the parabolic coordinates, which are defined as roots of the quadratic equation $\xi^2 + 2y\xi - x^2 = 0$.

4 Conclusions

In the previous paragraphs we have examined the six-parameters quasi-point canonical transformation leading to separation of variables for the Henon-Heiles system and a system with quartic potential. We have proved that this transformation strictly connected to the rectangular separated equations and that it can not be generalized. In addition we found some relations of these variables of separation with the two-dimensional supersymmetric quantum mechanics.

This work was supported by Swedish Royal Academy of Science grant, project N1314 for collaboration with Russian Academy of Science, and RFBR grant.

References

- [1] A.A. Andrianov, M.V. Joffe, and D.N. Nishnianidze. *Phys.Lett.*, A201:103–107, 1995. *Zap.Nauchn.Seminars POMI*, 224:68–80, 1995.
- [2] S. Baker, V.Z. Enolskii, and A. P. Fordy. *Phys.Lett*, A201:167–174, 1995.
- [3] M. Blaszak and S. Rauch-Wojciechowski. *J.Math.Phys.*, 35:1693–1709, 1994.
- [4] L.D. Landau and E.M. Lifshitz. *Mechanics*. Pergamon, London, 1960.
P.M. Morse and H. Feshbach. *Methods of theoretical physics*. McGraw-Hill, New York, 1953.
- [5] I. Marshall and S. Wojciechowski. *J.Math.Phys.*, 29:1338–1346, 1988.
- [6] A. Ramani, B. Dorizzi, B. Grammaticos, and J. Hietarinta. *Physica*, D18:171–179, 1986.
- [7] V. Ravozon, L. Gavrilov, and R. Cobo. *J.Math.Phys.*, 34:2385–2393, 1993.
- [8] V. Ravozon, A. Ramani, and R. Grammaticos. *Phys.Lett*, 191:91–95, 1994.
- [9] F.J. Romeiras. *J.Phys.*, A28:5633–5642, 1995.
- [10] E.T. Whittaker. *Analytical dynamics*. Cambridge University Press, Cambridge, 1973. §§152 and 153.
L.P. Eisenhart. *Phys.Rev.*, 45:427, 1934; 74:87, 1948.
- [11] M. Wojciechowska and S. Wojciechowski. *Phys.Lett.*, A 105:11–14, 1984.